

[illegible]

**ABSTRACT**

an acid addition salt of a mor-  
anic acid is provided.

ted from:

C<sub>16</sub>-dicarboxylic acids with mono-  
hols,  
ids,  
acids, selected from the group of  
kyl, hydroxyalkyl, alkoxyalkyl  
tuted benzoic acid, as well as of  
benzoic acids, which may option-  
the N atom,  
substituted 5-ring or 6-ring  
ing at least one N or S atom and  
oup function,  
ated, optionally substituted, oxo-  
ing 5 to 10 C atoms,  
r phenoxy-substituted saturated  
acids.

or heterocyclic C<sub>2</sub>- to C<sub>12</sub>-amino ac-  
no group is substituted with an -  
ed - C<sub>2</sub>- to C<sub>6</sub>-alkanoyl group or  
tituted - benzoyl group.

**ABSTRACT**

an acid addition salt of a mor-  
anic acid is provided.

ted from:

C<sub>16</sub>-dicarboxylic acids with mono-  
hols,  
ids,  
acids, selected from the group of  
kyl, hydroxyalkyl, alkoxyalkyl  
tuted benzoic acid, as well as of  
benzoic acids, which may option-  
the N atom,  
substituted 5-ring or 6-ring  
ing at least one N or S atom and  
oup function,  
ated, optionally substituted, oxo-  
ing 5 to 10 C atoms,  
r phenoxy-substituted saturated  
acids.

or heterocyclic C<sub>2</sub>- to C<sub>12</sub>-amino ac-  
no group is substituted with an -  
ed - C<sub>2</sub>- to C<sub>6</sub>-alkanoyl group or  
tituted - benzoyl group.

- ABSTRACT**
- an acid addition salt of a mor-  
anic acid is provided.
- ted from:
- C<sub>16</sub>-dicarboxylic acids with mono-  
hols,  
ids,  
acids, selected from the group of  
kyl, hydroxyalkyl, alkoxyalkyl  
tuted benzoic acid, as well as of  
benzoic acids, which may option-  
the N atom,  
substituted 5-ring or 6-ring  
ing at least one N or S atom and  
oup function,  
ated, optionally substituted, oxo-  
ing 5 to 10 C atoms,  
r phenoxy-substituted saturated  
acids.
- or heterocyclic C<sub>2</sub>- to C<sub>12</sub>-amino ac-  
no group is substituted with an -  
ed - C<sub>2</sub>- to C<sub>6</sub>-alkanoyl group or  
tituted - benzoyl group.